Table S2. Characters and character states used to generate the matrix of morphological

| Characters | States | | | | |
|---|-------------|-------------|---------|-------|--------|
| | 0 | 1 | 2 | 3 | 4 |
| 1- yellow cubital stripes on dorsal hindwings, | Very narrow | Narrow | Medium | Broad | Absent |
| when present | | | | | |
| 2- two series of doubled intervenal submarginal | Absent | Present | - | - | - |
| yellow spots, the internal series with spots larger | | | | | |
| than the external series | | | | | |
| 3- first line of the doubled intervenal | Weak | Strong | - | - | - |
| submarginal yellow spots on dorsal hindwings | through the | through the | | | |
| | end | end | | | |
| 4- ventral hindwing with red cubital spots | Absent | Present | - | - | - |
| 5- yellow streak over the forewing cubitus | Absent | Present | Usually | - | - |
| | | | Margin | | |
| 6- tip of the yellow streak over the hindwing | Thin | Rounded | - | - | - |
| cubitus | | | | | |
| 7- postmedian red band on dorsal forewing | Broad | Narrow | - | - | - |
| 8- proximity of postmedian red band on dorsal | Near | Distant | - | - | - |
| forewing to the margin | | | | | |
| 9-yellow cubital stripes on dorsal hindwing | Present | Absent | | - | - |
| mixed with numerous black scales, resulting in a | | | | | |
| grayish tone | | | | | |
| 10-Tone of the transverse red postmedian band | Orangish | Red | - | - | - |

characters of wing color patterns for *H. hermathena* subspecies.